arXiv:1606.04496v1 [math.GN] 14 Jun 2016

Effros, Baire, Steinhaus and Non-Separability

By A. J. Ostaszewski

Abstract. We give a short proof of an improved version of the Effros Open Mapping Principle via a shift-compactness theorem (also with a short proof), involving 'sequential analysis' rather than separability, deducing it from the Baire property in a general Baire-space setting (rather than under topological completeness). It is applicable to absolutely-analytic normed groups (which include complete metrizable topological groups), and via a Steinhaus-type Sum-set Theorem (also a consequence of the shift-compactness theorem) includes the classical Open Mapping Theorem (separable or otherwise).

Keywords: Open Mapping Theorem, absolutely analytic sets, base-σ-discrete maps, demi-open maps, Baire spaces, Baire property, group-action shift-compactness.

Classification Numbers: 26A03; 04A15; 02K20.

1 Introduction

We generalize a classic theorem of Effros [Eff] beyond its usual separable context. Viewed, despite the separability, as a group-action counterpart of the Open Mapping Theorem OMT (that a surjective continuous linear map between Fréchet spaces is open – cf. [Rud]), it has come to be called the *Open Mapping Principle* – see [Anc, §1]. Our 'non-separable' approach is motivated by a sequential property related to the Steinhaus-type Sum-set Theorem (that 0 is an interior point of $A-A$, for non-meagre A with BP, the Baire property –[Oxt], [Pic]), because of the following argument (which goes back to Pettis [Pet]).

Consider $L : E \to F$, a linear, continuous surjection between Fréchet spaces, and U a neighbourhood (nhd) of the origin. Choose A an *open* nhd of the origin with $A-A \subseteq U$; as $L(A)$ is non-meagre (since $\{nL(A) : n \in \mathbb{N}\}$ covers F) and has BP (see Proposition 2 in §2.3), $L(A) - L(A)$ is a nhd of the origin by the Sum-set Theorem. But of course

$$L(U) \supseteq L(A) - L(A),$$

so $L(U)$ is a nhd of the origin. So L is an open mapping.[1]

Throughout this paper, without further comment, all spaces considered will be metrizable, but not necessarily separable. We recall the Birkhoff-Kakutani theorem (cf. [HewR, §II.8.3]), that a metrizable group G with neutral element e_G has a right-invariant metric d_R^G. Passage to $||g|| := d_R^G(g, e_G)$ yields a (group) norm (invariant under inversion, satisfying the triangle inequality, cf. [ArhT], [BinO1] – see §4.1), which justifies calling these *normed groups*; any Fréchet space qua additive group, equipped with an F-norm ([KalPR, Ch. 1 §2]), is a natural example (cf. *Auth* in §2.2). Recall that a *Baire space* is one in which Baire's theorem holds – see [AaL]. Below we need the following.

Definitions 1 (cf. [Pet]). For G a metrizable group, say that $\varphi : G \times X \to X$ is a *Nikodym* group action (or that it has the Nikodym property – cf. [BinO4]) if for every non-empty open neighbourhood U of e_G and every $x \in X$ the set $Ux = \varphi_x(U) := \varphi(x, U)$ contains a non-meagre *Baire set*. (Here Baire set, as opposed to Baire space as above, means 'set with the Baire property'.)

2. A^q denotes the *quasi-interior* of A – the largest open set U with $U \backslash A$ meagre (cf. [Ost1, §4]); other terms ('analytic', 'base-σ-discrete', 'group action') are recalled later.

Concerning when the above property holds see §2.3. Our main results are Theorems S and E below, with Corollaries in §2.3 including OMT; see below for commentary.

Theorem S (Shift-compactness Theorem). *For T a Baire non-meagre subset of a metric space X and G a group, Baire under a right-invariant metric, and with separately continuous and transitive Nikodym action on X:*

for every convergent sequence x_n with limit x and any Baire non-meagre $A \subseteq G$ with $e_G \in A^q$ and $A^q x \cap T^q \neq \emptyset$, there are $\alpha \in A$ and an integer N such that $\alpha x \in T$ and

$$\{\alpha(x_n) : n > N\} \subseteq T.$$

In particular, this is so if G is analytic and all point-evaluation maps φ_x are base-σ-discrete.

[1] This proof is presumably well-known – so simple and similar to that for the automatic continuity of homomorphisms – but we have no textbook reference; cf. [KalPR, Cor. 1.5].

This theorem has wide-ranging consequences, including Steinhaus' Sum-set Theorem – see the survey article [Ost4], and the recent [BinO3]; cf. §4.2-4.

Theorem E (Effros Theorem – Baire version). *If*
(i) *the normed group G has separately continuous and transitive Nikodym action on X;*
(ii) *G is Baire under the norm topology and X is non-meagre*
– then for any open neighbourhood U of e_G and any $x \in X$ the set $Ux := \{u(x) : u \in U\}$ is a neighbourhood of x, so that in particular the point-evaluation maps $g \to g(x)$ are open for each x. That is, the action of G is micro-transitive.

In particular, this holds if G is analytic and Baire, and all point-evaluation maps φ_x are base-σ-discrete.

By Proposition B2 (§2.3) X, being non-meagre here, is also a Baire space.

The classical counterpart of Theorem E has G a Polish group; van Mill's version [vMil1] requires the group G to be analytic (i.e. the continuous image of some Polish space, cf. [JayR], [Kec2]). The Baire version above improves the version given in [Ost3], where the group is almost complete. (The two cited sources taken together cover the literature.)

A result due to Loy [Loy] and to Hoffmann-Jørgensen [HofJ, Th. 2.3.6 p. 355] asserts that a Baire, separable, analytic *topological group* is Polish (as a consequence of an analytic group being metrizable – for which see again [HofJ, Th. 2.3.6]), so in the analytic separable case Theorem E reduces to its classical version.

Unlike the proof of the Effros Theorem attributed to Becker in [Kec1, Th. 3.1], the one offered here does not employ the Kuratowski-Ulam Theorem (the Category version of the Fubini Theorem), a result known to fail beyond the separable context (as shown in [Pol], cf. [vMilP], but see [FreNR]).

For recent work on the circumstances when Theorems E and S are equivalent, see [BinO4]. For further commentary (connections between convexity and the Baire property, relation to van Mill's separation property in [vMil2], certain specializations) see §4.

2 Analyticity, micro-action, shift-compactness

We recall some definitions from general topology, before turning to ones that are group-related. We refer to [Eng] for general topological usage (but prefer

'meagre' to 'of first category'); see also §4.5,6.

2.1 Analyticity

We say that a subspace S of a metric space X has a *Souslin-$\mathcal{H}$ representation* if there is a *determining system* $\langle H(i|n)\rangle := \langle H(i|n) : i \in \mathbb{N}^{\mathbb{N}}\rangle$ of sets in $\mathcal{H}$ with ([Rog], [Han1], [Han2])

$$S = \bigcup_{i\in I} \bigcap_{n\in\mathbb{N}} H(i|n),\ (I := \mathbb{N}^{\mathbb{N}},\ i|n := (i_1, ..., i_n)).$$

A topological space is an (absolutely) *analytic* space if it is embeddable as a Souslin-$\mathcal{F}$ set in its own metric completion (with $\mathcal{F}$ the closed sets); in particular, in a complete metric space $\mathcal{G}_\delta$-subsets (being $\mathcal{F}_{\sigma\delta}$) are analytic. For more recent generalizations see e.g. [NamP]. According to Nikodym's theorem, if $\mathcal{H}$ above comprises Baire sets, then also S is Baire (the Baire property is preserved by the Souslin operation): so analytic subspaces are Baire sets. For background – see [Kec2] Th. 21.6 (the Lusin-Sierpiński Theorem) and the closely related Cor. 29.14 (Nikodym Theorem), cf. the treatment in [Kur] Cor. 1 p. 482, or [JayR] pp. 42-43. For the extended Souslin operation of non-separable descriptive theory see also [Ost2]. This motivates our interest in analyticity as a carrier of the Baire property, especially as continuous images of separable analytic sets are separable, hence Baire.

However, the continuous image of an analytic space is not in general analytic – for an example of failure see [Han3] Ex. 3.12. But this does happen when, additionally, the continuous map is base-σ-discrete, as defined below (*Hansell's Theorem*, [Han3] Cor. 4.2)**.** This technical condition is the standard assumption for preservation of analyticity and holds automatically in the separable realm. Special cases include *closed surjective* maps and *open-to-analytic injective* maps (taking open sets to analytic sets). To define the key concept just mentioned, recall that for an (indexed) family $\mathcal{B} := \{B_t : t \in T\}$:
(i) $\mathcal{B}$ is *index-discrete* in the space X (or just *discrete* when the index set T is understood) if every point in X has a nhd meeting the sets B_t for at most one $t \in T$,
(ii) $\mathcal{B}$ is *σ-discrete* if $\mathcal{B} = \bigcup_n \mathcal{B}_n$ where each set $\mathcal{B}_n$ is discrete as in (i), and
(iii) $\mathcal{B}$ is a *base for* $\mathcal{A}$ if every member of $\mathcal{A}$ is the union of a subfamily of $\mathcal{B}$.
For $\mathcal{T}$ a topology (the family of all open sets) with $\mathcal{B} \subseteq \mathcal{T}$ a base for $\mathcal{T}$, this reduces to $\mathcal{B}$ being simply a (topological) *base*.

Definitions. 1. ([Mic1], Def. 2.1) Call $f : X \to Y$ *base-σ-discrete* (or *co-σ-discrete,* [Han3, §3]) if the image under f of any discrete family in X has a σ-discrete base in Y.
2 ([Han3, §2], cf. [KanP]). An indexed family $\mathcal{A} := \{A_t : t \in T\}$ is *σ-discretely decomposable* if there are discrete families $\mathcal{A}_n := \{A_{tn} : t \in T\}$ such that $A_t = \bigcup_n A_{tn}$ for each t.
3 ([Mic1], Def. 3.3). Call $f : X \to Y$ *index-σ-discrete* if the image under f of any discrete family $\mathcal{E}$ in X is σ-discretely decomposable in Y. (Note $f(\mathcal{E})$ is regarded as indexed by $\mathcal{E}$, so could be discrete without being index-discrete.)

2.2 Action, micro-action, shift-compactness

Recall that a normed group G *acts continuously* on X if there is a continuous mapping $\varphi : G \times X \to X$ such that $\varphi(e_G, x) = x$ and $\varphi(gh, x) = \varphi(g, \varphi(h, x))$ $(x \in X, g, h \in G)$.The action φ is *separately continuous* if $g : x \mapsto \varphi(g, x)$ is continuous for each g, and $\varphi_x : g \mapsto \varphi(g, x)$ is continuous for each x; in such circumstances:
(i) the elements $g \in G$ yield autohomeomorphisms of X via $g : x \mapsto g(x) := \varphi(g, x)$ (as g^{-1} is continuous), and
(ii) point-evaluation of these homeomorphisms, $\varphi_x(g) = g(x)$, is continuous.
In certain situations joint continuity of action is implied by separate continuity (see [Bou] and literature cited in [Ost2]).

The action is *transitive* if for any x, y in X there is $g \in G$ such that $g(x) = y$. For later purposes (§2.3 and 3), say that the action of G on X is *weakly micro-transitive* if for $x \in X$ and each nhd A of e_G the set

$$\mathrm{cl}(Ax) = \mathrm{cl}\{ax : a \in A\}$$

has x as an interior point (in X). The action is *micro-transitive* ('transitive in the small' – for details see [vMil1]) if for $x \in X$ and each nhd A of e_G the set

$$Ax = \{ax : a \in A\}$$

is a nhd of x. This (norm) property implies that Ux is open for U open in G (i.e. that here each φ_x is an open mapping). We refer to Ax as an x *orbit* (the A-orbit of x). The following group action connects the Open Mapping Theorem to the present context.

Example (Induced homomorphic action). A surjective, continuous homomorphism $\lambda : G \to H$ between normed groups induces a transitive

action of G on H via $\varphi^{\lambda}(g,h) := \lambda(g)h$ (cf. [Ost2] Th. 5.1), specializing for G, H Fréchet spaces (regarded as normed, additive groups) and $\lambda = L : G \to H$ linear (Ancel [Anc] and van Mill [vMil1]) to

$$\varphi^{L}(a,b) := L(a) + b.$$

Of course for Fréchet spaces, by the Open Mapping Theorem itself, φ^{L} has the Nikodym property.

Definitions. 1. $Auth(X)$ denotes the autohomeomorphisms of a metric space (X, d^X); this is a group under composition. $\mathcal{H}(X)$ comprises those $h \in Auth(X)$ of bounded norm:

$$||h|| := \sup_{x \in X} d^X(h(x), x) < \infty.$$

2. For a normed group G acting on X, say that X has the *crimping property* (property C for short) w.r.t. G if, for each $x \in X$ and each sequence $\{x_n\} \to x$, there exists in G a sequence $\{g_n\} \to e_G$ with $g_n(x) = x_n$. (This and a variant occurs in [Ban, Ch. III; Th.4]; and [ChCh]; for the term see [BinO2].)

For a subgroup $\mathcal{G} \subseteq \mathcal{H}(X)$, say that X has the *crimping property* w.r.t. $\mathcal{G}$ if X has the crimping property w.r.t. to the natural action $(g,x) \to g(x)$ from $\mathcal{G} \times X \to X$. (This action is continuous relative to the left or right norm topology on $\mathcal{G}$ – cf. [Dug] XII.8.3, p. 271.)

3. As a matter of convenience, say that the *Effros property* (or *property E*) holds for the group G acting on X if the action is micro-transitive, as above.

4. For a subgroup $\mathcal{G} \subseteq Auth(X)$ say that X is *$\mathcal{G}$-shift-compact* (or, shift-compact under $\mathcal{G}$) if for any convergent sequence $x_n \to x_0$, any open subset U in X and any Baire set T co-meagre in U, there is $g \in \mathcal{G}$ with $g(x_n) \in T \cap U$ along a subsequence. Call the space *shift-compact* if it is $\mathcal{H}(X)$-shift-compact (cf. [MilO], [Ost5]).

In such a space, any Baire non-meagre set is locally co-meagre (co-meagre on open sets) in view of Prop. B2 below.

We shall prove in § 3.1 equivalence between the Effros and Crimping properties:

Theorem EC. *The Effros property holds for a group G acting on X iff X has the Crimping property w.r.t. G.*

We now clarify the role of shift-compactness.

Proposition B1. *For any subgroup $\mathcal{G} \subseteq \mathcal{H}(X)$, if X is $\mathcal{G}$-shift-compact, then X is a Baire space.*

Proof. We argue as in [vMil2] Prop 3.1 (1). Suppose otherwise; then X contains a non-empty meagre open set. By Banach's Category Theorem (or localization principle, for which see [JayR] p. 42, or [Kel] Th. 6.35), the union of all such sets is a largest open meagre set M, and is non-empty. Thus $X\backslash M$ is a co-meagre Baire set. For any $x \in M$ the constant sequence $x_n \equiv x$ is convergent and, since $X\backslash M$ is co-meagre in X, there is $g \in G$ with $g(x) \in X\backslash M$. But, as g is a homeomorphism, $g(M)$ is a non-empty open meagre set, so is contained in M, implying $g(x) \in M$, a contradiction. □

A similar argument gives the following and clarifies an assumption in Theorem E.

Proposition B2 (cf. [vMil2]; [HofJ, Prop. 2.2.3]). *If X is non-meagre and G acts transitively on X, then X is a Baire space.*

Proof. As above, refer again to M, the union of all meagre open sets, which, being meagre, has non-empty complement. For x_0 in this complement and any non-empty open U pick $u \in U$ and $g \in G$ such that $g(x_0) = u$. Now, as g is continuous, $g^{-1}(U)$ is a nhd of x_0, so is non-meagre, since every nhd of x_0 is non-meagre. But g is a homeomorphism, so $U = g(g^{-1}(U)$ is non-meagre. So X is Baire, as every non-empty open set is non-meagre. □

2.3 Nikodym actions

The following result generalizes one that, for separable groups G, is usually a first step in proving the weakly micro-transitive variant of the classical Effros Theorem (cf. Ancel [Anc] Lemma 3, [Ost3] Th. 2). Indeed, one may think of it as giving a form of 'very weak micro-transitivity'.

Proposition 1. *If G is a normed group, acting transitively on a non-meagre space X with each point evaluation map $\varphi_x : g \mapsto g(x)$ base-σ-discrete relative – then for each non-empty open U in G and each $x \in X$ the set Ux is non-meagre in X.*

In particular, if G is analytic, then G is a Nikodym action.

Proof. We first work in the right norm topology, i.e. derived from the assumed right-invariant metric $d_R^G(s,t) = ||st^{-1}||$. Suppose that $u \in U$, and

so without loss of generality assume that $U = B_\varepsilon(u) = B_\varepsilon(e_G)u$ (open balls of radius some $\varepsilon > 0$); then put $y := ux$ and $W = B_\varepsilon(e_G)$. Then $Ux = Wy$. Next work in the left norm topology, derived from $d^G_L(s,t) = ||s^{-1}t|| = d^G_R(s^{-1}, t^{-1})$ (for which $W = B_\varepsilon(e_G)$ is still a nhd of e_G). As each set hW for $h \in G$ is now open (since now the left shift $g \to hg$ is a homeomorphism), the open family $\mathcal{W} = \{gW : g \in G\}$ covers G. As G is metrizable (and so has a σ-discrete base), the cover $\mathcal{W}$ has a σ-discrete refinement, say $\mathcal{V} = \bigcup_{n\in\mathbb{N}} \mathcal{V}_n$, with each $\mathcal{V}_n$ discrete. Put $X_n := \bigcup\{Vy : V \in \mathcal{V}_n\}$; then $X = \bigcup_{n\in\mathbb{N}} X_n$, as $X = Gy$, and so X_n is non-meagre for some n, for $n = N$ say. Since φ_y is base-σ-discrete, $\{Vy : V \in \mathcal{V}_N\}$ has a σ-discrete base, say $\mathcal{B} = \bigcup_{m\in\mathbb{N}} \mathcal{B}_m$, with each $\mathcal{B}_m$ discrete. Then, as $\mathcal{B}$ is a base for $\{Vy : V \in \mathcal{V}_N\}$,

$$X_N = \bigcup\nolimits_{m\in\mathbb{N}} \left(\bigcup\{B \in \mathcal{B}_m : (\exists V \in \mathcal{V}_N) B \subseteq Vy\} \right).$$

So for some m, say for $m = M$,

$$\bigcup\{B \in \mathcal{B}_m : (\exists V \in \mathcal{V}_N) B \subseteq Vy\}$$

is non-meagre. But as $\mathcal{B}_M$ is discrete, by Banach's Category Theorem (cf. Prop. B1), there are $\hat{B} \in \mathcal{B}_M$ and $\hat{V} \in \mathcal{V}_N$ with $\hat{B} \subseteq \hat{V}y$ such that $\hat{B}$ is non-meagre. As $\mathcal{V}$ refines $\mathcal{W}$, there is some $\hat{g} \in G$ with $\hat{V} \subseteq \hat{g}W$, so $\hat{B} \subseteq \hat{V}y \subseteq \hat{g}Wy$, and so $\hat{g}Wy$ is non-meagre. As $\hat{g}^{-1}$ is a homeomorphism of X, $Wy = Ux$ is also non-meagre in X.

If G is analytic, then as U is open, it is also analytic (since open sets are $\mathcal{F}_\sigma$ and Souslin-$\mathcal{F}$ subsets of analytic sets are analytic, cf. [JayR]), and hence so is $\varphi_x(U)$. Indeed, since φ_x is continuous and base-σ-discrete, Ax is analytic (Hansell's Theorem, §2.1), so Souslin-$\mathcal{F}$, and so Baire by Nikodym's Theorem (§2.1). □

Definition. **(**Ancel [Anc]**).** Call the map φ_x *countably-covered* if there exist self-homeomorphisms h^x_n of X for $n \in \mathbb{N}$ such that for any open nhd U in G the sets $\{h^x_n(\varphi_x(U)) : n \in \mathbb{N}\}$ cover X.

Proposition 1′ (cf. Ancell [Anc]) *For the action $\varphi : G \times X \to X$ with X non-meagre, if each map φ_x is countably-covered and takes open sets to sets with the Baire property, then the action has the Nikodym property.*

Proof. If φ_x is countably-covered, then there exist self-homeomorphisms h^x_n of X for $n \in \mathbb{N}$ such that for any open nhd U in G the sets $\{h^x_n(\varphi_x(U)) :$

$n \in \mathbb{N}\}$ cover X. Then for X non-meagre, there is $n \in \mathbb{N}$ with $h_n^x(\varphi_x(U))$ non-meagre, so $Ux = \varphi_x(U)$ is itself non-meagre, being a homeomorphic copy of $h_n^x(\varphi_x(U))$. As Ux is assumed Baire, the action has the Nikodym property. $\square$

For E separable, an immediate consequence of *continuous* maps taking open sets to analytic sets (which are Baire sets) and of Prop. $1'$ is that φ^L is a Nikodym action.

For the general context, one needs *demi-open* continuous maps, which preserve almost completeness (absolute $\mathcal{G}_\delta$ sets modulo meagre sets – see [Mic2] and its antecedent [Nol]), as it is not known which linear maps are base-σ-discrete – a delicate matter to determine, since the former include continuous linear surjections (by Lemma 1 below) and preserve almost analyticity as opposed to analyticity.

For present purposes, however, the *monotonicity property* below suffices. We omit the proof of the following observation (for which see the opening step in [Rud, 2.11], or [Con, Ch. 3 §12.3], or the Appendix). For the underlying translation-invariant metric of a Fréchet space denote below by $B(a, r)$ the open r-ball with centre a.

Lemma 1. *For a continuous linear map $L : X \to Y$ from a Fréchet space X to a normed space Y, for $s < t < r$*

$$\mathrm{int}(\mathrm{cl}L(B(0, s))) \subseteq L(B(0, t)) \subseteq L(B(0, r)).$$

Hence for $L(a, r)$ convex, either $L(B(a, r))$ is meagre or differs from $\mathrm{int}L(B(a, r))$ by a meagre set.

Proposition 2. *For L a continuous linear surjection from a Fréchet space E to a non-meagre normed space F, the action φ^L has the Nikodym property.*

Proof. As in Prop $1'$ for $L : E \to F$ a continuous linear surjection, $\{\varphi_x^L : x \in F\}$ are countably-covered. Indeed, fixing $x \in F$

$$h_n^x(z) := n(z - x) \qquad (n \in \mathbb{N} \text{ and } z \in F)$$

is on the one hand a self-homeomorphism satisfying $h_n^x(\varphi_x(L(V))) = L(nV)$, since $n[(L(v) + x) - x] = nL(v) = L(nv)$; on the other hand the family

$$\{h_n^x(L(V) + x) : n \geq 1\}$$

covers F, as $\{nV : n \in \mathbb{N}\}$ covers E for V any open nhd of the origin in E (by the 'absorbing' property, cf. [Con, 4.1.13], [Rud, 1.33]). In particular, $nL(B(0,1))$ is non-meagre for some n, and so $L(B(0,s))$ is non-meagre for any s. By Lemma 1, $L(B(0,t))$ for any $t > s$ contains the non-meagre Baire set $\mathrm{cl}L(B(0,s))$. □

Corollary 1 below is now immediate; it is used in [Ost2, Th. 5.1] to prove the 'Semi-Completeness Theorem', an Ellis-type theorem [Ell, Cor. 2] (cf. [Ost6]) giving a one-sided continuity condition which implies that a right-topological group generated by a right-invariant metric is a topological group.

Corollary 1 (cf. [Ost2, Th. 5.1], 'Open Homomorphism Theorem'). *If the continuous surjective homomorphism λ between normed groups G and H, with G analytic and H a Baire space, is base-σ-discrete, then λ is open; in particular, for λ bijective, λ^{-1} is continuous.*

Corollary 2. *For $L : E \to F$ a continuous surjective linear map between Fréchet spaces, the point evaluations φ_b^L for $b \in F$ are open, and so L is an open mapping.*

Proof. By surjectivity of L, the action is transitive, and by Prop 2 the action φ^L has the Nikodym property. So by Theorem E above the point-evaluations maps φ_b^L are open. Hence so also is L. □

3 Proofs

3.1 Proof that E $\Longleftrightarrow$ C

In [BinO1] Th. 3.15 we showed that if the Effros property holds for the action of a group G on X, then X has the crimping property w.r.t. G. We recall the argument, as it is short. Suppose that $x = \lim x_n$. For each n, take $U = B^G_{1/n}(e_G)$; then $Ux := \{u(x) : u \in U\}$ is an open nhd of x, and so there exists $h_{n,m} \in U$ with $h_{n,m}(x) = x_m$ for all m large enough, say for all $m > m(n)$. Without loss of generality we may assume that $m(1) < m(2) < ....$. Put $h_m := e_G$ for $m < m(1)$, and for $m(k) \leq m < m(k+1)$ take $h_m := h_{k,m}$. Then $h_m \in B^G_{1/k}(e_G)$, so h_m converges to e_G and $h_m(e_G) = x_m$.

For the converse, suppose that the Effros property fails for G acting on X. Then for some open nhd U of e_G and some $x \in X$, $Ux := \{u(x) : u \in U\}$ is not an open nhd of x. So for each n there is a point $x_n \in B_{1/n}(x) \backslash Ux$. As x_n converges to x there are homeomorphisms h_n converging to the identity e_G with $h_n(x) = x_n$. As U is an open nhd of e_G and since h_n converges to e_G, there is N such that $h_n \in U$ for $n > N$. In particular, for any $n > N$, $h_n(x) = x_n \in Ux$, a contradiction.

3.2 Weak S

We view Th. S as having 'two tasks': to find a 'translator of the sequence' τ, and to locate it in a given Baire non-meagre subset of the group – provided that subset satisfies a consistency condition (a necessary condition).

For clarity we break the tasks the into two steps – the first delivering a weaker version of S in Proposition 3 below. The arguments are based on the following lemma. We note a corollary, observed earlier by van Mill in the case of metric topological groups ([vMil2, Prop. 3.4]), which concerns a co-meagre set, but we need its refinement to a localized version for a non-meagre set.

Separation Lemma. *Let G be a normed group, with separately continuous and transitive Nikodym action on a non-meagre space X. Then for any point x and any F closed nowhere dense, $W_{x,F} := \{\alpha \in G : \alpha(x) \notin F\}$ is dense open in G. In particular, G separates points from nowhere dense closed sets.*

Proof. The set $W_{x,F}$ is open, being of the form $\varphi_x^{-1}(X \backslash F)$ with φ_x continuous (by assumption). By the Nikodym property, for U any non-empty open set in G, the set Ux is non-meagre, and so $Ux \backslash F$ is non-empty, as F is meagre. But then for some $u \in U$ we have $u(x) \notin F$. $\square$

Corollary 2. *If G is a normed group, Baire in the norm topology with transitive and separately continuously Nikodym action on a non-meagre space X space, and T is co-meagre in X– then for countable $D \subseteq X$, the set $\{g : g(D) \subseteq T\}$ is a dense $\mathcal{G}_\delta$.*

In particular, this holds if G is analytic and each point-evaluation map $\varphi_x : g \to g(x)$ is base-σ-discrete.

Proof. Without loss of generality, the co-meagre set is of the form $T = U \backslash \bigcup_{n \in \omega} F_n$ with each F_n closed and nowhere dense, and U open. Then, by

the Separation Lemma and as G is Baire,

$$\{g \in G : g(D) \subseteq T\} = \bigcap\nolimits_{n\in\omega}\{g : g(D) \cap F_n = \emptyset\} = \bigcap\nolimits_{d\in D, n\in\omega}\{g : g(d) \notin F_n\}$$

is a dense $\mathcal{G}_\delta$. □

Proposition 3. *If T is a Baire non-meagre subset of a metric space X and G a normed group, Baire in its norm topology, acting separately continuously and transitively on X, with the Nikodym property – then, for every convergent sequence x_n with limit x_0 there is $\tau \in G$ and an integer N with $\tau x_0 \in T$ and*

$$\{\tau(x_n) : n > N\} \subseteq T.$$

Proof. Write $T := M \cup (U \setminus \bigcup_{n\in\omega} F_n)$ with U open, M meagre and each F_n closed and nowhere dense in X. Let $u_0 \in T \cap U$. By transitivity there is $\sigma \in G$ with $\sigma x_0 = u_0$. Put $u_n := \sigma x_n$. Then $u_n \to u_0$. Put

$$C := \bigcap\nolimits_{m,n\in\omega}\{\alpha \in G : \alpha(u_m) \notin F_n\},$$

a dense $\mathcal{G}_\delta$ in G; then, by the Separation Lemma above, as G is Baire,

$$\{\alpha \in G : \alpha(u_0) \in U\} \cap C$$

is non-empty. For α in this set we have $\alpha(u_0) \in U \setminus \bigcup_{n\in\omega} F_n$. Now $\alpha(u_n) \to \alpha(u_0)$, by continuity of α, and U is open. So for some N we have for $n > N$ that $\alpha(u_n) \in U$. Since $\{\alpha(u_m) : m = 1, 2, ..\} \in X \setminus \bigcup_{n\in\omega} F_n$, we have for $n > N$ that $\alpha(u_n) \in U \setminus \bigcup_{n\in\omega} F_n \subseteq T$.

Finally put $\tau := \alpha\sigma$; then $\tau(x_0) = \alpha\sigma(x_0) \in T$ and $\{\tau(x_n) : n > N\} \subseteq T$. □

3.3 Proof of S

We work in the right norm topology and use the notation of the preceding proof (of Proposition 3), so that U here is the quasi-interior of T and $\sigma x_0 = u_0$. As $e_G \in A^q$ and A is a non-meagre Baire set, we may without loss of generality write $A = B_\varepsilon(e_G) \setminus \bigcup_n G_n$, where each G_n is closed nowhere dense with $e_G \notin G_n$ and $B_\varepsilon(e_G)$ is the quasi-interior of A.

As $A^q x_0 \cap T^q$ is non-empty, there is $\alpha_0 \in B_\varepsilon(e_G)$ with $\alpha_0 x_0 \in U$ (but, we want a better α so that $\alpha x_0 \in T$ and $\alpha \in A$). Put $\beta_0 = \alpha_0\sigma^{-1}$; then

$$\begin{aligned}\beta_0 &= \alpha_0\sigma^{-1} \in B_\varepsilon(e_G)\sigma^{-1} \cap \{\alpha : \alpha(x_0) \in U\}\sigma^{-1} \\ &= B_\varepsilon(e_G)\sigma^{-1} \cap \{\beta : \beta(\sigma x_0) \in U\} = B_\varepsilon(e_G)\sigma^{-1} \cap \{\beta : \beta(u_0) \in U\},\end{aligned}$$

i.e. the open set $\{\beta : \beta(u_0) \in U\} \cap B_\varepsilon(e_G)\sigma^{-1}$ is non-empty. So

$$(C \setminus \bigcup\nolimits_n G_n\sigma^{-1}) \cap \{\beta : \beta(u_0) \in U\} \cap B_\varepsilon(e_G)\sigma^{-1} \neq \emptyset,$$

since G is a Baire space and each $G_n\sigma^{-1}$ is closed and nowhere dense in G (as the right shift $g \to g\sigma^{-1}$ is a homeomorphism).

So there is β with $\beta(u_0) \in U$ such that $\alpha := \beta\sigma \in B_\varepsilon(e_G) \setminus \bigcup_n G_n = A$. That is, $\alpha x_0 = \beta u_0 \in U$; so $\beta(u_n) \in U$ for large n, for $n > N$ say, as $\alpha x_0 = \lim \alpha x_n = \lim \beta\sigma x_n = \lim \beta u_n$. But $\{\beta(u_m) : m = 1, 2, ..\} \in X \setminus \bigcup_n F_n$, as $\beta \in C$; so $\beta(u_n) \in U \setminus \bigcup_n F_n \subseteq T$ for $n > N$.

Finally, $\alpha(x_0) = \beta\sigma(x_0) \in T$ and $\{\alpha(x_n) : n > N\} \subseteq T$. □

3.4 Proof that S $\Longrightarrow$ E

Assume G acts transitively on X and that X is non-meagre. Let $B := B_\varepsilon(e_G)$ and suppose that for some x the set Bx is not a nhd of x. Then there is $x_n \to x$ with $x_n \notin Bx$ for each n. Take $A := B_{\varepsilon/2}(e_G)$ and note first that A is a symmetric open set ($A^{-1} = A$, since $||g|| = ||g^{-1}||$), and secondly that by the Nikodym property Ax contains a non-meagre, Baire subset T. So by Theorem S, as Ax meets T^q, there are $a \in A$ (which being open has the Baire property) and a co-finite $\mathbb{M}_a$ such that $ax_m \in Ax$ for $m \in \mathbb{M}_a$. For any such m, choose $b_m \in A$ with $ax_m = b_m x$. Then $x_m = a^{-1}b_m x \in A^2x \subseteq Bx$, a contradiction (note that $a^{-1} \in A$, by symmetry).

As earlier, in the special case that G is (metrizable and) analytic, A is analytic, since open sets are $\mathcal{F}_\sigma$ and Souslin-$\mathcal{F}$ subsets of analytic sets are analytic, cf. [JayR, Th. 2.5.3], by Prop. 3 Ax is Baire non-meagre, as φ_x is base-σ-discrete.

4 Complements

1. Normed groups. For T an *algebraic* group with neutral element $e = e_G$, say that $||\cdot|| : T \to \mathbb{R}_+$ is a *group-norm* ([ArhT], [BinO1]) if the following properties hold:

(i) *Subadditivity* (Triangle inequality): $||st|| \leq ||s|| + ||t||$;

(ii) *Positivity*: $||t|| > 0$ for $t \neq e$ and $||e|| = 0$;

(iii) *Inversion* (Symmetry): $||t^{-1}|| = ||t||$.

Then $(T, ||.||)$ is called a *normed-group.*

The group-norm generates a right and a left *norm topology* via the right-invariant and left-invariant metrics $d_R^T(s,t) := ||st^{-1}||$ and $d_L^T(s,t) := ||s^{-1}t|| = d_R^T(s^{-1},t^{-1})$. In the right norm-topology the right shift $\rho_t(s) := st$ is a uniformly continuous homeomorphism, since $d_R(sy,ty) = d_R(s,t)$, so in particular the group is a right topological group; likewise in the left norm-topology the left shift $\lambda_s(t) = st$ is a uniformly continuous homeomorphism. Since $d_L^T(t,e) = d_L^T(e,t^{-1}) = d_R^T(e,t)$, convergence at e is identical under either topology. One may refer to whichever topology is appropriate, since despite their differences, they are homeomorphic. Thus one may say, for instance, that separability is a *norm property*, as separability of either norm topology implies that of the other.

See [Ost5] for a characterization of almost-complete normed groups as non-meagre almost-analytic (i.e. analytic modulo a meagre set).

2. Genericity Our focus on various shift-compactness theorems derives from their close affinity with the literature of 'generic' automorphisms (for which see [Mac], also described in the introduction to [Ost5]) and from a multitude of its applications, for which see e.g. [BinO1], and these offering a unifying sequential compactness-like, or combinatorial, perspective on category-measure duality and on other, apparently unrelated, problems.

3. Assumptions in Theorem S. With regard to the assumption of separate continuity, note that a theorem of Bouziad ([Bou, Th. 3]) implies that a separately continuous action by a metrizable left-topological Baire group acting on a metric space is in fact jointly continuous; we retain the only apparently weaker hypothesis of separate continuity, because there are variants of Theorem S, where joint continuity is absent (and the group is not Baire): see [MilO]. The result is connected with van Mill's *Separation Property*: say that X has SP ([vMil2]) with respect to a group of homeomorphisms $\mathcal{G}$ if for any countable set D in X and any meagre set M in X, there is a homeomorphism $g \in \mathcal{G}$ such that $g(D) \cap M = \emptyset$; it is interesting to recall here the Galvin-Mycielski-Solovay characterization of subsets of $\mathbb{R}$ of strong measure zero as sets A such that for any meagre M there is g with $(g + A) \cap M = \emptyset$ (see [Mil, §3]). Stated equivalently, SP asserts that for D countable and T co-meagre, there is $g \in \mathcal{G}$ with $g(D) \subseteq T$ (see also the next remark). Compared to this restatement, Theorem S refers on the one hand to a smaller class of countable sets (convergent sequences, or their co-finite parts), but on the other hand asserts embeddability into a larger class of sets – sets T that

are 'locally' rather than globally co-meagre; for further information see also [MilO].

4. Shift-compactness and the SP property. In view of the strength of Th. S and to place our results in context, we briefly summarize some of the relevant results of [vMil2, Th. 1.1]. A (separable metric) space with SP is Baire, by the proof which we imported for Propositions B1 and B2 in §2; likewise, an almost complete non-meagre space with SP is completely Baire. Since an absolutely co-analytic space is Polish iff it is 'completely Baire', i.e. closed-hereditarily Baire (closed subspaces are Baire), for which see Kechris [Kec2, Cor. 21.21], it follows that an absolutely Borel space with the SP is Polish ([vMil2, Th. 1.1]). More generally, if an analytic group G acts on a space X and SP holds w.r.t. G, then X is Polish. Van Mill also shows from his Prop 3.4 (cf. our Prop. 2) that a locally compact homogeneous space has the SP. It seems likely that, just as with Proposition B, more of these arguments can be copied across in the language of Th. S.

It is noted in [BinO2] that a Polish space which is strongly locally homogeneous has SP.

5. *Index-σ-discrete maps.* In many circumstances it is easier to work with index-σ-discrete maps: see [Ost1] for a brief discussion of this point and for relations with the automatic continuity of homomorphisms (noted earlier in [Nol]). Furthermore, if $f : X \to Y$ is injective and *closed-analytic,* i.e. carries closed sets to analytic sets (or, alternatively *open-analytic*, mutatis mutandis), then f is base-σ-discrete – in fact index-σ-discrete ([Han3] Prop. 3.14). On the other hand, if f is surjective and closed with Y metrizable, then f is base-σ-discrete ([Han3] Prop. 3.10). So if the group action is such that each φ_x is open-analytic and *injective* (as when a vector space acts on itself), then each φ_x is index- and so base-σ-discrete.

6. *Generalizations of analyticity.* A Hausdorff space is *almost analytic* if it is analytic modulo a meagre set. Similarly, a space X' is absolutely $\mathcal{G}_\delta$, or an *absolute-$\mathcal{G}_\delta$*, if X' is a $\mathcal{G}_\delta$ in all spaces X containing X' as a subspace. This latter property is equivalent to complete metrizability in the realm of metrizable spaces [Eng, Th. 4.3.24] (and to topological/Čech completeness in the realm of completely regular spaces – [Eng, §3.9]). So a metrizable absolute-$\mathcal{G}_\delta$ is analytic. A metric space is *almost complete* if it contains a dense absolute $\mathcal{G}_\delta$. The notion of 'almost completeness' is due to Frolík in [Frol] (but its name to Michael [Mic2] – see also [AaL] and [BinO1]).

Consequently, it is natural to assume that when a group G acts on a space X each point evaluation map $\varphi_x : g \to g(x)$ is not only continuous but also base-σ-discrete; the *Nikodym property* above is implied by the former property when G is analytic. Note that [Han3] Ex. 3.12 shows that, for $D = (\mathbb{R}, d_{\text{discrete}})$ the discrete space of cardinality the continuum, the projection from $D \times [0,1] \to [0,1]$ is an open mapping that is not base-σ-discrete (by reference to the closed discrete graph of a bijection between D and $[0,1]$).

Index-σ-discrete maps (easier to work with) are base-σ-discrete ([Han3] Prop. 3.7(i)); the latter combined with continuity preserves analyticity, as mentioned earlier. Evidently, if all evaluation maps φ_x are continuous and base-σ-discrete and A is an open subset of an analytic group G, acting on X, then A is analytic, and so $\varphi_x(A) = Ax$ is analytic; thus the point evaluations are then open-analytic. So our Nikodym hypothesis on the action is not far from demanding that point evaluations be open-analytic.

7. *Convexity and Baire Property.* We have seen above that analyticity confers the Baire property. Convexity may also confer it: for A convex, if $\text{int}A$, its interior, is non-empty, then $\text{int}A$ is dense in A, and so A has the Baire property: $A \backslash \text{int}(A) \subseteq \text{cl}(A) \backslash \text{int}(\text{cl}(A))$, a nowhere dense set; so A differs from its interior by a nowhere dense subset of $\text{cl}(A) \backslash \text{int}(\text{cl}(A))$. If A is non-meagre its closure has non-empty interior, raising the question of whether A itself has non-empty interior.

The Banach context is particularly transparent; Banach's lemma (see Appendix) yields: *for a continuous linear map $T : X \to Y$ from a Banach space X to a normed space Y and $x \in X$, if $B(0,\rho) \subseteq \text{cl}(T(\bar{B}(x,r)))$ with $r, \rho > 0$, then $B(0,\rho) \subseteq T(\bar{B}(x,r))$.*

So, as above, by translation, interpret this lemma as asserting that for $r > 0$ and $A := L(\bar{B}(0,r))$ one has $\text{int}(\text{cl}(A)) \subseteq \text{int}(A)$. So $A \backslash \text{int}(A) \subseteq \text{cl}(A) \backslash \text{int}(\text{cl}(A))$, a nowhere dense set, so A differs from its interior by a nowhere dense subset of $\text{cl}(A) \backslash \text{int}(\text{cl}(A))$, and so has the Baire property.

In the Fréchet case, with the balls referring to the underlying translation-invariant metric, one may apply Lemma 1 of §2.3. Taking the convex set $B(0,r)$ with $r > 0$, suppose $T(B(0,r))$ is non-meagre; then $T(B(0,s))$ is non-meagre for some $0 < s < r$. By the claim, for $s < t < r$,

$$\emptyset \neq \text{int}(\text{cl}T(B(0,s))) \subseteq T(B(0,t)).$$

As $T(B(0,t))$ is convex, $\text{int}T(B(0,t))$ is convex, and being non-empty is dense in $\text{cl}T(B(0,t))$, i.e. $\text{cl}T(B(0,t)) = \text{cl}(\text{int}T(B(0,t)))$. So $T(B(0,t))$

differs from $\mathrm{int}T(B(0,t))$ by a meagre set, as

$$\begin{aligned} T(B(0,t))\backslash \mathrm{int}T(B(0,t)) &\subseteq \mathrm{cl}T(B(0,t))\backslash \mathrm{int}T(B(0,t)) \\ &= \mathrm{cl}(\mathrm{int}T(B(0,t)))\backslash \mathrm{int}T(B(0,t)). \end{aligned}$$

Finally, taking $s < t_n \nearrow r$, one has

$$T(B(0,r))\backslash \mathrm{int}T(B(0,r)) \subseteq \bigcup\nolimits_{n\in\mathbb{N}} \mathrm{cl}(\mathrm{int}T(B(0,t_n)))\backslash \mathrm{int}T(B(0,t_n)).$$

So for $B(0,r)$ convex, *either* $T(B(0,r))$ *is meagre or differs from* $\mathrm{int}T(B(0,r))$ *by a meagre set.*

8. *Topological and semitopological groups.* When a group G equipped with a topology such that the (action) map $(g_1, g_2) \to g_1 g_2$ from G^2 to G is separately continuous (i.e. left and right shifts are continous), then it is said to be a *semitopological group*. Thus a topological group is semitopological. If also the topology is metrizable and G is Baire, Theorem S applies and asserts here that if T is non-meagre in G and $g_n \to g$, then for some $t \in T$ the point tg is in T and almost all of the sequence tg_n is in T. As in Theorem S, under these additional assumptions, group multiplication in G is jointly continuous; indeed, Bouziad [Bou] proves that a semitopological Baire p-space (and metric spaces are p-spaces) has jointly continuous multiplication (is 'paratopological').

9. Specializing the proof of Theorem S to semitopological groups. The argument proving Th. S is particularly transparent in the case of a semitopological group and follows a standard real-line argument, as follows. Suppose $x_n \to x_0$ and consider $T := U\backslash \bigcup F_m$ with F_m closed and nowhere dense, and U open. For any $u_0 \in T \cap U$, take $\sigma(x) = x x_0^{-1} u_0$, which is a continuous right-shift, so that $u_n = \sigma(x_n) = x_n(x_0^{-1}u_0) \to u_0 = \sigma(x_0)$. Now

$$u_0^{-1}U \cap \bigcap\nolimits_{n,m} u_n^{-1}(X\backslash F_m)$$

is a dense $\mathcal{G}_\delta$, since left-shifts are homeomorphisms. As G is Baire, there is g with

$$g \in u_0^{-1}U \cap \bigcap\nolimits_{n,m} u_n^{-1}(X\backslash F_m).$$

Then $u_0 g \in U$ and $u_0 g \notin F_m$ for all m, so $u_0 g \in T$. By continuity of right-shifts, $u_n g \to u_0 g$, so for large n, say for $n > N$, we have

$$u_n g \in U.$$

For each such n we have $u_n g \notin F_m$ for all m and so $u_n g \in T$. Thus, as $u_n g = x_n(x_0^{-1}u_0)g$, we conclude the existence for some $g \in G$ and $u_0 \in T$ with $t := u_0 g \in T$ that, as in Theorem S

$$x_n x_0^{-1} t \in T \text{ for } n > N.$$

10. From E back to S. In Proposition 3 density of the set $W_{x,F}$ was deduced from a Lemma which, as noted, asserts a very weak microtransitivity. Unsurprisingly, Property E also implies density of that set (see below). However, Property E does not imply that G is Baire, since (see Introduction) there do exist meagre analytic groups acting on a non-meagre space which by van Mill's result have Property E notwithstanding. So we can go no further with the density argument of Proposition 3. Of course, as in Proposition 3, if we know that G is Baire, then E implies S.

As for the density claim, consider $\beta \in G$. Suppose first that $\beta(x) \in F$. By the Effros property, the set $B_\varepsilon(\beta)x = B_\varepsilon(e_G)\beta x$ is an open nhd of βx. As F is nowhere dense and closed, there is $y \in \text{int}(B_\varepsilon(e_G)\beta x)\backslash F$. So there is $\gamma \in B_\varepsilon(e_G)$ such that $y = \alpha x := \gamma\beta x \notin F$. So $a = \gamma\beta \in B_\varepsilon(\beta) \cap \{\alpha : \alpha(x) \notin F\}$.

If, on the other hand, $\beta(x) \notin F$, then $\beta \in B_\varepsilon(\beta) \cap \{\alpha : \alpha(x) \notin F\}$. This proves the claim.

A personal note. Whilst the present author's entry into mathematics owes hugely both to Karol Borsuk and Ambrose Rogers (thesis advisor), being confirmed as a topologist is down to a first meeting with Mary Ellen Rudin in 1972 (at the Keszthely conference) and subsequent frequent stays at UW Madison, visiting her and the wonderous set-theoretic community there. It is thus a pleasure to dedicate this paper especially to her memory and to express once more the great debt to UW friends and colleagues, among whom also was Anatole Beck, recently passed away.

Acknowledgement. I thank the Referee for some thought-provoking comments that influenced the final presentation, and Henryk Toruńczyk for drawing my attention to Ancel's work and related literature.

References

[AaL] J. M. Aarts and D. J. Lutzer, Completeness properties designed for recognizing Baire spaces, *Dissertationes Math.* **116** (1974), 48pp.

[Anc] F. D. Ancel, An alternative proof and applications of a theorem of E. G. Effros, *Michigan Math. J.* **34** (1987), 39–55.

[ArhT] A. Arhangelskii, M. Tkachenko, *Topological groups and related structures*. Atlantis Studies in Mathematics, 1. Atlantis Press; World Scientific, 2008.

[Ban] S. Banach, *Théorie des opérations linéaires*, Monografie Mat. **1**, 1932 (in: "Oeuvres", Vol. 2, PWN, 1979.), translated as '*Theory of linear operations*', North Holland, 1987.

[BinO1] N. H. Bingham, A. J. Ostaszewski, Normed versus topological groups: dichotomy and duality, *Dissertationes Math.* **472** (2010), 138 pp.

[BinO2] N. H. Bingham, A. J. Ostaszewski,Topological regular variation: I Slowly varying functions, *Topology Appl.*. **157** (2010), 1999-2013.

[BinO3] N. H. Bingham, A. J. Ostaszewski, The Steinhaus theorem and regular variation: de Bruijn and after, *Indagationes Math.* **24** (2013), 679-692.

[BinO4] N. H. Bingham and A. J. Ostaszewski, Beyond Lebesgue and Baire IV: Density topologies and a converse Steinhaus-Weil theorem, preprint.

[Bou] A. Bouziad, Continuity of separately continuous group actions in p-spaces, *Topology Appl.* **71** (1966), 119-124.

[ChCh] J. J. Charatonik and W. J. Charatonik, The Effros metric, *Topology Appl.* **110** (2001), 237-255.

[Con] J. B. Conway, *A course in functional analysis.* 2nd ed. Graduate Texts in Mathematics **96**, Springer, 1990.

[Dug] J. Dugundji, *Topology,* Allyn and Bacon, 1966.

[Eff] E. G. Effros, Transformation groups and C*-algebras. *Ann. of Math.* (2) **81** (1965), 38–55.

[Ell] R. Ellis, Continuity and homeomorphism groups. *Proc. Amer. Math. Soc.* **4**, (1953) 969–973.

[Eng] R. Engelking, *General Topology,* Heldermann Verlag, Berlin 1989.

[FreNR] D. Fremlin, T. Natkaniec, I. Recław, Universally Kuratowski-Ulam spaces, *Fund. Math.* **165** (2000), 239–247.

[Frol] Z. Frolík, Generalizations of the G_δ-property of complete metric spaces, *Czechoslovak Math. J.* **10** (85) (1960), 359–379.

[Han1] R. W. Hansell, On the nonseparable theory of Borel and Souslin sets, *Bull. American Math. Soc.* **78.2** (1972), 236-241.

[Han2] R. W. Hansell, On the representation of nonseparable analytic sets, *Proc. American Math. Soc.* **39.2** (1973), 402-408.

[Han3] R. W. Hansell, On characterizing non-separable analytic and extended Borel sets as types of continuous images, *Proc. London Math. Soc.* (3) **28** (1974), 683-699.

[HewR] E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis, I: Structure of topological groups, integration theory, group representations*. Grundl. math. Wiss. **115**, Springer, 1963.

[HofJ] J. Hoffmann-Jørgensen, Automatic continuity, Section 3 of [TopHJ].

[JayR] J. E. Jayne, C. A. Rogers, Analytic sets, Part 1 of [Rog]

[KalPR] N. J. Kalton, N. T. Peck, J. R. Roberts, *An F-space sampler*, LMS Lect. Notes Ser. 89, CUP, 1984.

[KanP] J. Kaniewski and R. Pol , Borel-measurable selectors for compact-valued mappings in the non-separable case, *Bull. Acad. Polon. Sci.* **23** (1975), 1043-1050.

[Kec1] A. S. Kechris, Topology and descriptive set theory, *Topology Appl.* **58** (1994), no. 3, 195–222.

[Kec2] A. S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics 156, Springer, 1995.

[Kel] J. L. Kelley, *General Topology*, Van Nostrand, 1955.

[Kur] K. Kuratowski, *Topology,* Vol. I., PWN, Warsaw 1966.

[Loy] R. J. Loy, Multilinear mappings and Banach algebras. *J. London Math. Soc.* (2) **14.3** (1976), 423–429.

[Mac] D. Macpherson, A survey of homogeneous structures, *Discrete Math.* **311** (2011), 1599-1634.

[Mic1] E. Michael, On maps related to σ-locally finite and σ-discrete collections of sets*, Pacific J. Math.* **98** (1982) 139-152.

[Mic2] E. Michael, Almost complete spaces, hypercomplete spaces and related mapping theorems, *Topology Appl.* **41** (1991), 113–130.

[vMil1] J. van Mill, A note on the Effros Theorem, *Amer. Math. Monthly* **111.9** (2004), 801-806.

[vMil2] J. van Mill, Analytic groups and pushing small sets apart, *Trans. Amer. Math. Soc.* **361** (2009), 5417–5434.

[vMilP] J. van Mill, R. Pol, The Baire category theorem in products of linear spaces and topological groups.*Topology Appl.* **22** (1986), no. 3, 267–282.

[Mil] A. Miller, Special subsets of the real line, Ch. 5 in *Handbook of Set-theoretic topology*, ed. K. Kunen and J. E. Vaughan, Elsevier, 1984, 201-233.

[MilO] H. I. Miller and A. J. Ostaszewski, Group-action and Shift-compactness, *J. Math. Analysis and App.* **392** (2012), 23-39.

[NamP] I. Namioka, R. Pol, σ-fragmentability and analyticity. *Mathematika* **43** (1969), 172-181.

[Nol] D. Noll, A topological completeness concept with applications to the open mapping theorem and the separation of convex sets. *Topology Appl.* **35** (1990), 53-69.

[Oxt] J. C. Oxtoby, *Measure and category, a survey of the analogies between topological and measure spaces*, 2nd ed., Grad. Texts Math. 2, Springer, 1980 (1st ed. 1971).

[Ost1] A. J. Ostaszewski, Analytically heavy spaces: Analytic Cantor and Analytic Baire spaces, *Topology Appl.* **158** (2011), 253-275.

[Ost2] A. J. Ostaszewski, Analytic Baire spaces, *Fund. Math.*, **217** (2012), 189-210.

[Ost3] A. J. Ostaszewski, Almost completeness and the Effros Open Mapping Principle in normed groups, *Topology Proceedings*, **41** (2013), 99-110.

[Ost4] A. J. Ostaszewski, Shift-compactness in almost analytic submetrizable Baire groups and spaces, *Topology Proceedings*, **41** (2013), 123-151.

[Ost5] A. J. Ostaszewski, Beyond Lebesgue and Baire III: Steinhaus' Theorem and its descendants, *Topology Appl.* **160** (2013), 1144-1154.

[Ost6] A. J. Ostaszewski, The semi-Polish Theorem: One-sided vs joint continuity in groups, *Topology Appl.* **160** (2013), 1155-1163

[Pet] B. J. Pettis, On continuity and openness of homomorphisms in topological groups. *Ann. of Math. (2)* **52** (1950). 293–308.

[Pic] S. Piccard, Sur les ensembles de distances des ensembles de points d'un espace Euclidien. *Mém. Univ. Neuchâtel* **13**, Neuchâtel, 1939.

[Pol] R. Pol, Note on category in Cartesian products of metrizable spaces, *Fund. Math.* **102** (1979), 55–59.

[Rog] C. A. Rogers, J. Jayne, C. Dellacherie, F. Topsøe, J. Hoffmann-Jørgensen, D. A. Martin, A. S. Kechris, A. H. Stone, *Analytic sets*, Academic Press, 1980.

[Rud] W. Rudin, *Functional Analysis*, McGraw-Hill, 2nd ed. 1991 (1st ed. 1973).

[TayL] A. E. Taylor and D. C. Lay, *Introduction to Functional Analysis*, Wiley, 1980 (1st ed., A. E. Taylor, Wiley, 1958).

[TopHJ] F. Topsøe, J. Hoffmann-Jørgensen, Analytic spaces and their applications, Part 3 of [Rog].

Mathematics Department, London School of Economics, Houghton Street, London WC2A 2AE
a.j.ostaszewski@lse.ac.uk

Appendix

For convenience we include here the

Lemma. (see e.g. [TayL, Lemma 5.4]; [Ban, Ch. III; Proof of Th.3]) *For a continuous linear map $T : X \to Y$ from a Banach space X to a normed space Y, if $B(0,\rho) \subseteq \mathrm{cl}(T(\bar{B}(a,r)))$, then $B(0,\rho) \subseteq T(\bar{B}(a,r))$.*

Proof. Let ε be arbitrary with $0 < \varepsilon < 1$ and suppose that $B(0,\rho) \subseteq \mathrm{cl}(T(\bar{B}(a,r)))$. If $y \in B(0,\rho)$, then $||y - y_1|| < \varepsilon\rho$ for some $y_1 = Tx_1$ with $x_1 \in B(a,r)$. So $x_1 = a + z_1$ with $||z_1|| < r$. But $||y - Tx_1|| < \varepsilon\rho$ and, by homogeneity of the norm in Y,

$$B(0,\varepsilon\rho) = \varepsilon B(0,\rho) \subseteq \mathrm{cl}T(\bar{B}(\varepsilon a, \varepsilon r)),$$

so

$$||y - Tx_1 - Tx_2|| < \varepsilon^2\rho.$$

for some $x_2 \in \bar{B}(\varepsilon a, \varepsilon r)$, i.e. $x_2 = \varepsilon a + z_2$ for some z_2 with $||z_2|| \leq \varepsilon r$. Continue by induction. Now

$$x_1 + ... + x_n = a(1 + \varepsilon + \varepsilon^2 + ... + \varepsilon^{n-1}) + z_1 + .. + z_n,$$

and

$$||z_1 + ... + z_n|| \leq ||z_1|| + ... + ||z_n|| \leq r(1 + ... + \varepsilon^{n-1}).$$

By completeness, $x_1 + ... + x_n \to x_* \in \bar{B}(a/(1-\varepsilon), r/(1-\varepsilon)$ and $Tx_* = y$. So

$$B(0,\rho) \subseteq T(\bar{B}(a/(1-\varepsilon), r/(1-\varepsilon))$$

and so

$$B(0,(1-\varepsilon)\rho) = (1-\varepsilon)B(0,\rho) \subseteq T(\bar{B}(a,r)).$$

As ε was arbitrary we conclude that

$$B(0,\rho) = \bigcup\nolimits_{0<\varepsilon<1} B(0,(1-\varepsilon)\rho) \subseteq T(\bar{B}(a,r)).$$

In particular $0 \in \mathrm{int}(T(\bar{B}(a,r)))$. □